

Acupuncture's Role in Solving the Opioid Epidemic:

Evidence, Cost-Effectiveness, and Care Availability for Acupuncture as a Primary, Non-Pharmacologic Method for Pain Relief and Management

White Paper 2017

**The American Society of Acupuncturists (ASA)
The American Alliance for Professional Acupuncture Safety (AAPAS)
The Acupuncture Now Foundation (ANF)
The American TCM Association (ATCMA)
The American TCM Society (ATCMS)
National Federation of TCM Organizations (NFTCMO)**

Original authors: The Joint Acupuncture Opioid Task Force:

Chair: Bonnie M. Abel Bolash, MAc, LAc.

Member organizations:

The Acupuncture Now Foundation (ANF)

The American Society of Acupuncturists (ASA)

**Original Contributors: Matthew Bauer, LAc
Bonnie Bolash, LAc
Lindy Camardella, LAc
Mel Hopper Koppelman, MSc
John McDonald, PhD, FAACMA
Lindsay Meade, LAc
David W Miller, MD, LAc**

First Revising Author: Arthur Yin Fan, CMD, PhD, LAc (ATCMA)

Corresponding Author: David W Miller, MD, LAc

**Revising Authors: Sarah Faggert, DAc, LAc
Hongjian He, CMD, LAc
Mel Hopper Koppelman, MSc
Yong Ming Li, MD, PhD, LAc
Amy Matecki, MD, LAc*
David W Miller, MD, LAc
John Pang, MD**
Jun Xu, MD, LAc**

***Division Chief, Dept. of Medicine, Highland Hospital, Alameda Health System**

****Division of Otolaryngology-Head and Neck Surgery, Department of Surgery,
University of California, San Diego School of Medicine**

**Acupuncture's Role in Solving the Opioid Epidemic:
Evidence, Cost-Effectiveness, and Care Availability for Acupuncture
as a Primary, Non-Pharmacologic Method for Pain Relief and Management
White Paper 2017**

Abstract

The United States is facing a national opioid epidemic, and medical systems are in need of non-pharmacologic strategies that can be employed to decrease the public's opioid dependence. Acupuncture has emerged as a powerful, evidence-based, safe, cost-effective, and available treatment modality suitable to meeting this need. Acupuncture has been shown to be effective for the management of numerous types of pain, and mechanisms of action for acupuncture have been described and are understandable from biomedical, physiologic perspectives. Further, acupuncture's cost-effectiveness could dramatically decrease health care expenditures, both from the standpoint of treating acute pain and through avoiding the development of opioid addiction that requires costly care, destroys quality of life, and can lead to fatal overdose. Numerous federal regulatory agencies have advised or mandated that healthcare systems and providers offer non-pharmacologic treatment options, and acupuncture stands as the most evidence-based, immediately available choice to fulfil these calls. Acupuncture can safely, easily, and cost-effectively be incorporated into hospital settings as diverse as the emergency department, labor and delivery suites, and neonatal intensive care units to treat a variety of pain seen commonly in hospitals. Acupuncture is already being successfully and meaningfully utilized by the Veterans Administration and various branches of the U.S. Military.

1. Acupuncture is an effective, safe, and cost-effective treatment for numerous types of acute and chronic pain. Acupuncture should be recommended as a first line treatment for pain before opiates are prescribed, and may reduce opioid use.

1.1 Effectiveness/Efficacy of acupuncture for different types of pain.

There is growing research evidence to support the effectiveness and efficacy of acupuncture for the relief of numerous types of pain, especially chronic pain, as well as the use of acupuncture for a diverse array of conditions (Table 1¹, Appendix 1). Acupuncture has been shown to be effective for treating various types of pain, with the strongest evidence emerging for **back pain, neck pain, shoulder pain, chronic headache, and osteoarthritis**. In an individual patient meta-analysis of 17,922 people from 29 randomized controlled trials (RCTs), it was concluded that the effect sizes in comparison to no acupuncture controls were 0.55 standard deviation (SD), 95% confidence interval (CI) [0.51-0.58] for back and neck pain; 0.57 SD, 95% CI [0.50-0.64] for osteoarthritis; and 0.42 SD, 95% CI [0.37-0.46] for chronic headache. No meta-analysis was performed on shoulder pain as there were only three eligible RCTs. In all analyses, true acupuncture was significantly superior to no acupuncture and sham acupuncture controls ($p < 0.001$).²

Table 1. Acupuncture for the use of numerous conditions including pain conditions.	
The Acupuncture Evidence Project	
(Mar 2013 - Sept 2016)	
Evidence of positive effect	
<ul style="list-style-type: none"> ● Allergic rhinitis (perennial & seasonal) ● Chemotherapy-induced nausea and vomiting (CINV) (with anti-emetics) ● Chronic low back pain ● Headache (tension-type and chronic) ● Knee osteoarthritis ● Migraine prophylaxis ● Post-operative nausea & vomiting ● Post-operative pain 	
Evidence of potential positive effect	
<ul style="list-style-type: none"> ● Acute low back pain ● Acute stroke ● Ambulatory anaesthesia ● Anxiety ● Aromatase-inhibitor-induced arthralgia ● Asthma in adults ● Back or pelvic pain during pregnancy ● Cancer pain ● Cancer-related fatigue ● Constipation ● Craniotomy anaesthesia ● Depression (with antidepressants) ● Dry eye ● Hypertension (with medication) ● Insomnia ● Irritable bowel syndrome ● Labor pain ● Lateral elbow pain ● Menopausal hot flashes 	<ul style="list-style-type: none"> ● Modulating sensory perception thresholds ● Neck pain (some types/non-whiplash) ● Obesity ● Peri-menopausal & post-menopausal insomnia ● Plantar heel pain ● Post-stroke insomnia ● Post-stroke shoulder pain ● Post-stroke spasticity ● Post-traumatic stress disorder ● Prostatitis pain/chronic pelvic pain syndrome ● Recovery after colorectal cancer resection ● Restless leg syndrome ● Schizophrenia (with antipsychotics) ● Sciatica ● Shoulder impingement syndrome (early stage) (with exercise) ● Shoulder pain ● Smoking cessation (up to 3 months) ● Stroke rehabilitation ● Temporomandibular joint disorder

In the largest study of its kind to date, 454,920 patients were treated with acupuncture for headache, low back pain, and/or osteoarthritis in an open pragmatic trial. Effectiveness was rated by the 8,727 treating physicians as marked or moderate in 76% of cases.³

In a network meta-analysis comparing different physical interventions for pain from knee osteoarthritis, acupuncture was found to be superior to sham acupuncture, muscle-strengthening exercise, tai chi, weight loss, standard care, and aerobic exercise (in ranked order). Acupuncture was found to be statistically more significantly effective than muscle-strengthening exercises, standardised mean difference (SMD) = 0.49, 95% CI [0.00-0.98].⁴

In early 2017, the American College of Physicians (ACP) published guidelines based on the evidence for the non-invasive treatment of low back pain. For acute or subacute low back pain, the ACP strongly recommends non-pharmacologic treatment with **acupuncture**, along with superficial heat, massage, or spinal manipulation, and nonsteroidal anti-inflammatory drugs or skeletal muscle relaxants. For chronic low back pain, the ACP also strongly recommends **acupuncture**, in addition to exercise, multidisciplinary rehabilitation, mindfulness-based stress reduction, tai chi, yoga, motor control exercise, progressive relaxation, electromyography biofeedback, low-level laser therapy, operant therapy, cognitive behavioral therapy, and spinal manipulation, etc.⁵

A systematic review and meta-analysis on acupuncture for the treatment of **sciatica** concluded that acupuncture was superior to standard pharmaceutical care (such as ibuprofen, diclofenac, and prednisone) in reducing pain intensity (mean difference (MD) = 1.25, 95% CI [1.63-0.86]) and pain threshold (MD = 1.08, 95% CI [0.98–1.17]).⁶

A systematic review and network meta-analyses of 21 different interventions for sciatica found that acupuncture was second in global effect only to biological agents, and superior to all other interventions including non-opioid and opioid medications.⁷

A systematic review on acupuncture and moxibustion for **lateral elbow pain** found that acupuncture and moxibustion were superior or equal to standard care.⁸

A systematic review on acupuncture for **plantar heel pain** found that evidence supporting the effectiveness of acupuncture was comparable to the evidence available for standard care interventions such as stretching, night splints, and dexamethasone.⁹

The use of acupuncture to relieve **pain associated with surgical procedures** captured the world's attention in the early 1970's when well-known *New York Times* journalist James Reston, who, while in China, witnessed acupuncture's effectiveness on his post-operative pain, published his personal experience with acupuncture shortly before President Richard Nixon's trip to China. Since then, acupuncture has been used before, during, and after surgery to manage pain and to improve post-surgical recovery in a variety of contexts.^{10,11,12,13,14,15,16,17,18} It is noteworthy that acupuncture has been reported to be effective in pain relief during and after surgical procedures on children and animals as well.^{19,20,21,22}

Nonetheless, over the past two decades post-operative pain management has come to rely increasingly on opioids while underutilizing alternative analgesics such as acupuncture. In 2012, surgeons and dentists combined prescribed 16.2% of all opioids in the U.S, trailing only family practices as the leading source of opioid prescriptions at 18.2%.²³ Eighty to ninety-four percent of patients undergoing low risk surgical procedures fill a prescription for opioids within 7 days.^{24,25} Recent data has shown that opioid prescriptions vary widely and that the majority of surgical patients are over-prescribed opioids, as approximately 70% of pills go unused.²⁶ The risk of chronic

opioid use after surgery in previously non-dependent patients is determined to be 5.9 – 6.5%,²⁷ although in select populations such as head and neck cancer patients, the risk is up to 40%.²⁸ The increase in post-operative opioid use is somewhat paradoxical considering that known adverse effects such as sedation, pneumonia,^{29,30} ileus, urinary retention, and delirium prolong patient recovery and delay the meeting of discharge goals.³¹

Acupuncture is a promising adjunctive analgesic modality to reduce the risk of post-operative opioid dependence. A meta-analysis published in late 2017 in the *Journal of the American Medical Association (JAMA Surgery)* focused on non-pharmacological treatments in reducing pain after total knee arthroplasty. Thirty-nine randomized clinical trials were included in the meta-analysis (2,391 patients). Moderate-certainty level evidence showed that electrotherapy reduced the use of opioids (MD = 3.50, 95% CI [5.90-1.10]) morphine equivalents in milligrams per kilogram per 48 hours ($p = .004$, $I^2 = 17\%$), and that acupuncture delayed opioid use (MD = 46.17, 95% CI [20.84-71.50]) minutes to the first patient-controlled analgesia ($p < .001$, $I^2 = 19\%$). There was low-certainty level evidence that acupuncture improved pain (MD = 1.14, 95% CI [1.90-0.38] on a visual analog scale at 2 days ($p = .003$, $I^2 = 0\%$). Evidence showed that acupuncture is better than cryotherapy, continuous passive motion, and preoperative exercise in the studied condition.³² Reduction in opioid use has been demonstrated across a wide range of both minor and major surgical procedures, including cardiac surgery,³³ thoracic surgery,³⁴ and craniotomy.^{35,36} Moreover, acupuncture may even reduce post-operative ileus and expedite bowel recovery after colorectal cancer resection.³⁷ As acupuncture is often combined with electric stimulation, electro-acupuncture may have more clinical benefit in post-operative pain management.

A Cochrane systematic review on acupuncture or acupressure for **primary dysmenorrhea** found that both acupuncture and acupressure were more effective in reducing pain than placebo controls.³⁸ Five other systematic reviews and/or meta-analyses on various forms of acupoint stimulation including acupuncture, acupressure, and moxibustion for primary dysmenorrhea have reported similar outcomes.^{39,40,41,42,43}

The effectiveness of acupuncture for **labor pain** is still unclear, largely due to the heterogeneity of designs and methods in studies which have produced mixed results. While some studies reported no reduction in analgesic medications, some studies reported reduction of pain during labor, reduced use of opioid medications and epidural analgesia, and a shorter second stage of labor.^{44,45,46}

A systematic review on acupuncture for **trigeminal neuralgia** suggests that acupuncture may be equal or superior to carbamazepine, but the evidence is weakened by the low methodological quality of some included studies.⁴⁷

A Cochrane systematic review on acupuncture for **fibromyalgia** found low to moderate-level evidence that acupuncture improves pain and stiffness compared with no treatment and standard therapy. Furthermore, electro-acupuncture is probably better than manual acupuncture for pain in fibromyalgia, although more studies are warranted.⁴⁸

A prospective, randomized trial of acupuncture vs. morphine to treat **emergency department/emergency room patients with acute onset, moderate to severe pain** was conducted. Acupuncture provided more effective and faster analgesia than morphine, and was better tolerated. The study included 300 patients, with 150 patients in each group. Success rate was

significantly different between the 2 groups (92% in the acupuncture group vs 78% in the morphine group, $p < .001$). Resolution time was 16 ± 8 minutes in the acupuncture group vs 28 ± 14 minutes in the morphine group ($p < .005$). Overall, 89 patients (29.6%) experienced minor adverse effects; of these, 85 (56.6%) were in the morphine group and only 4 (2.6%) were in the acupuncture group ($p < .001$).⁴⁹

The above mentioned meta-analysis included 29 trials and 17,922 patients with chronic pain conditions; data on longer term follow-up (available for 20 trials, including 6376 patients) suggests that approximately 90% of the benefit of acupuncture relative to controls would be sustained at 12 months after the course of treatment. Patients can generally be reassured that treatment effects persist at least 12 months.⁵⁰

1.2 Safety and feasibility of acupuncture for pain management.

The strongest evidence for the safety of acupuncture in chronic pain management comes from an open pragmatic trial involving 454,920 patients who were treated for headache, low back pain, and/or osteoarthritis. Minor adverse events were reported in 7.9% of patients while only 0.003% (13 patients) experienced severe adverse events. Minor adverse events included needling pain, hematoma, and bleeding, while serious adverse events included pneumothorax, acute hyper- or hypotensive crisis, erysipelas, asthma attack, and aggravation of suicidal thoughts.⁵¹ In a prospective feasibility study, acupuncture was seen as feasible, safe, and acceptable in an ICU setting by patients from diverse backgrounds.⁵² A systematic review suggests that acupuncture performed by trained practitioners using clean needle technique is a generally safe procedure.⁵³ The medical literature also indicates that acupuncture may be used successfully on cancer patients for symptom management due to the low risks associated with its use.⁵⁴

1.3 Cost-effectiveness of acupuncture for pain management.

In a systematic review of 8 cost-utility and cost-effectiveness studies of acupuncture for chronic pain, the cost per quality adjusted life year (QALY) gained was below the thresholds used by the UK National Institute for Health and Clinical Excellence for “willingness to pay”. The chronic pain conditions included in the systematic review included low back pain, neck pain, dysmenorrhoea, migraine and headache, and osteoarthritis.⁵⁵ In a cost-effectiveness analysis of non-pharmacological treatments for osteoarthritis of the knee, acupuncture was found to be the most cost-effective option when analysis was limited to high-quality studies.⁵⁶ Using acupuncture for pain management, patients and insurers can save money and successfully manage their pain and other symptoms without the adverse risks associated with prescription medications. A recent study found that full insurance coverage for acupuncture would increase an average insured member’s monthly health insurance premium from \$0.38 to \$0.76. Acupuncture was noted to save \$35,480, \$32,000, \$9,000, and \$4,246 per patient for migraine, angina pectoris, severe osteoarthritis, and carpal tunnel syndrome respectively.⁵⁷ Considering the large fees associated with prescription medications and surgery for pain conditions, acupuncture is extremely cost-effective.

The Acupuncture Evidence Project also enumerates those conditions for which acupuncture has been found to be cost-effective (Table 2).⁵⁸

Table 2. Conditions with demonstrated evidence of cost effectiveness.

- Allergic rhinitis
- Low back pain
- Ambulatory anaesthesia
- Migraine
- Chronic pain: neck pain (plus usual medical care)
- Depression
- Osteoarthritis
- Dysmenorrhoea
- Post-operative nausea and vomiting
- Headache

1.4 Can adjunctive acupuncture treatment reduce the use of opioid-like medications?

Some studies have reported reduced consumption of opioid-like medication (OLM) by more than 60% following surgery when acupuncture is used.^{59,60} A pilot RCT also showed a reduction by 39% in OLM use in non-malignant pain after acupuncture, an effect which lasted fewer than 8 weeks after acupuncture treatment ceased.⁶¹ The above mentioned meta-analysis, having moderate-certainty level evidence, showed that electro-acupuncture therapy reduced the use of opioids, and acupuncture delayed opioid use; with low-certainty level evidence indicating that acupuncture improved pain.⁶² The conclusions suggest that electro-acupuncture may be effective in reducing or delaying the use of opioid medications.

In a study looking at acupuncture's effectiveness in treating pain in a military cohort at a United States Air Force medical center, acupuncture dramatically decreased the use of opiates and other pain medications among personnel. Opioid prescriptions decreased by 45%, muscle relaxants by 34%, NSAIDs by 42%, and benzodiazepines by 14%. Quality of life measures also showed impressive changes, with some measures of improvements showing statistical significance ($p < 0.001$).⁶³

The Veterans Administration is increasingly looking to incorporate acupuncture into care, as is the U.S. Air Force and other branches. Training of military physicians is increasing, and systems are being studied to further incorporate acupuncture. The military is rapidly incorporating this care into its offered services for service members.^{64,65}

Studies of the effects of opioid analgesia in the elderly reveal a significant burden of disease due to falls from mental impairment. This is worsened when seniors are using multiple medications affecting cognition. In a recent study from the Journal of the American Geriatric Society, serious falls as per Medicare Part A and B ICD/CPT codes were evaluated in 5,556 nursing home residents aged 65 or greater. Seniors taking > 3 + CNS standardized daily doses were more likely to have a serious fall than those not taking any CNS medications (adjusted odds ratio = 1.83, 95% CI [1.35-2.48]), and the authors urge, "Clinicians should be vigilant for opportunities to discontinue or decrease the doses of individual CNS medications and/or consider non-pharmacological alternatives."⁶⁶

A recent study in the New Zealand Medical Journal noted that medication related harms were both common and created a substantial burden of disease for patients and the healthcare system. They listed opioids first among the six categories of medications causing the most significant burden.⁶⁷ In light of the findings of these studies and similar, utilization of non-pharmacologic treatment options such as acupuncture must be a priority of paramount status.

2. Acupuncture's analgesic mechanisms have been extensively researched and acupuncture can increase the production and release of endogenous opioids in animals and humans.

Mechanisms underlying acupuncture's analgesic effects have been extensively researched for over 60 years. In animal models, acupuncture and/or electro-acupuncture has been shown to be effective for the alleviation of inflammatory, neuropathic, cancer-related, and visceral pain. Mechano-transduction of the needling stimulus at specific points on the body triggers the release of ATP and adenosine, which bind to local afferents.^{68,69} Ascending neural pathways involving A β , A δ , and C sensory fibres have been mapped, as well as a mesolimbic analgesic loop in the brain and brainstem, descending pathway mechanisms, dopaminergic contributors, and cytokine, glutamate, nitric oxide, and gamma-amino-butyric-acid (GABA) effects. Acupuncture analgesia has been shown to involve several classes of opioid neuropeptides including enkephalins, endorphins, dynorphins, endomorphins, and nociceptin (also known as Orphanin FQ). Among the non-opioid neuropeptides, substance P (SP), vasoactive intestinal peptide (VIP), and calcitonin gene-related peptide (CGRP) have been investigated for their roles in both the analgesic and anti-inflammatory effects of acupuncture.^{70,71,72,73}

Given that acupuncture analgesia activates the production and release of endogenous opioids and activates μ and δ opioid receptors, it is feasible that acupuncture, used in conjunction with OLM, might alleviate pain with a lower OLM dose for patients already taking OLM.⁷⁴ This idea is further supported by evidence that acupuncture increases μ opioid receptor binding potential, allowing for effective analgesia at lower doses of OLM.⁷⁵ For patients not yet prescribed OLM, acupuncture should be recommended prior to OLM prescription commencing. This would be in-line with existing guidelines, such as those by the ACP⁷⁶ and the Centers for Disease Control and Prevention (CDC),⁷⁷ which recommend that safe and effective non-opioid alternatives should first be exhausted before resorting to OLM.

It is important to note as well that opioids as a monotherapy are often not as successful as may be thought in the general public perception. A recent systematic review of opioid analgesics for low back pain, which included 7,925 participants, found that opioids were poorly tolerated and for those who tolerate them the effect is unlikely to be clinically important within guideline recommended doses.⁷⁸ The first ever randomized controlled trial evaluating the long-term effectiveness of opioids, found that those on long-term opioid analgesia were actually in marginally more pain at 12 months than those in the non-opioid group.⁷⁹ Hence, complementary methods of pain control are critical to successful patient management.

3. Acupuncture is effective for the treatment of chronic pain involving maladaptive neuroplasticity.

Adverse neuroplastic changes can present a challenge in pain management, as maladaptive neuroplasticity can be associated with severe chronic pain that is resistant to treatment. Via peripheral stimulation, acupuncture may relieve the symptoms of patients affected by problematic neuroplastic changes. There is evidence that acupuncture has the capacity to reverse adverse neuroplastic changes in the dorsal horns of the spine, as well as in the somatosensory cortex.^{80,81,82,83} This suggests that acupuncture may have an important role in treating chronic pain which involves adverse neuroplastic changes.

4. Acupuncture is a useful adjunctive therapy in opiate dependency and rehabilitation.

Acupuncture is an effective way to treat opioid addiction. There are more than 45 human and animal studies and clinical trials included in the PubMed database exploring acupuncture's role in minimizing the usage of multiple drugs of potential abuse including opiates and methamphetamine.⁸⁴ In 1973, Doctors Wen and Cheung, et al. from Hong Kong published an accidental finding that ear acupuncture treatment for respiratory patients had apparently alleviated opioid withdrawal signs and symptoms.⁸⁵ These findings were replicated by others around the world, including in New York and Sydney in the mid-1970s. In 1985, Dr. Michael Smith, et al. in New York established the National Acupuncture Detoxification Association (NADA), which today operates in over 40 countries with an estimated 25,000 providers. There are more than 1,000 programs in the U.S. and Canada that now use acupuncture to help addicts overcome their addictions.⁸⁶ Acupuncture for addiction is a versatile modality that can be effortlessly integrated into many environments including prisons, in- and outpatient programs, community centers, disaster relief, and humanitarian aid efforts. Furthermore, acupuncture addiction protocols can address acute and prolonged withdrawal symptoms, stress and anxiety related to drug withdrawal, and help prevent relapse. Using drugs to treat those already drug addicted is not a rational plan of action, and finding sound, non-pharmacologic treatment options is of paramount importance.

A meta-analysis done in 2012 concluded that “the majority [of studies] agreed on the efficacy of acupuncture as a strategy for the treatment of opiate addiction” and that “neurochemical and behavioral evidence have shown that acupuncture helps reduce the effects of positive and negative reinforcement involved in opiate addiction by modulating mesolimbic dopamine neurons. Moreover, several brain neurotransmitter systems involving opioids and GABA have been implicated in the modulation of dopamine release by acupuncture.”⁸⁷ In a recent RCT involving 28 newborns with Neonatal Abstinence Syndrome, laser acupuncture plus OLM significantly reduced the duration of oral morphine therapy when compared to OLM alone.⁸⁸ The mechanism for acupuncture in opiate withdrawal was found to be mediated by the endogenous opioid “dynorphin” binding to kappa-opioid receptors.⁸⁹

5. Acupuncture has been recommended as a first line non-pharmacologic therapy by the FDA, as well as the National Academies of Sciences, Engineering, and Medicine in coping with the opioid crisis. The Joint Commission has also mandated that hospitals provide non-pharmacologic pain treatment modalities.

The Food and Drug Administration (FDA) released proposed changes to its opioid prescription guidelines in early May 2017. This was titled as its “Blueprint for Prescriber Education for Extended-Release and Long-Acting Opioids”. The guidelines now recommend that doctors become informed about non-pharmacologic options for pain control to help avoid the overuse of opioids.⁹⁰ Per the FDA’s request, the National Academies of Sciences, Engineering, and Medicine (NASEM) released a report to outline the state of the science regarding prescription opioid abuse and misuse, as well as the evolving role that opioids play in pain management. The new NASEM report on pain management and opioids recommends more public education, reimbursement models, and support for non-drug approaches to pain treatment. It systematically summarizes the evidence for acupuncture’s clinical benefits in treating different pain conditions, and provides an overview of some of the basic science underlying acupuncture’s mechanisms in pain management.⁹¹ Further, effective January 1, 2018, the Joint Commission has mandated that hospitals provide non-pharmacologic pain treatment modalities.⁹² Acupuncture is ideally suited to fulfil this mandate. These official, evidence-based clinical guidelines are in line with global healthcare trends; as of November 2015, acupuncture had over 870 recommendations in official clinical guidelines for over 100 conditions from institutions in over 30 countries.⁹³

6. Among most non-pharmacological managements for pain relief now available, acupuncture therapy is the most effective and specific for opioid abuse and overuse.

There are several forms of non-pharmacological managements for acute and chronic pain, including physical therapy, spinal cord manipulation, yoga, tai chi, and cognitive behavioral therapy. Among these therapies commonly recommended by medical authorities, acupuncture is the most specific in targeting the endogenous opioid system. There is more evidence that acupuncture can induce endorphins to cope with acute and chronic pain in basic research than for any other non-pharmacological approach for pain.^{94,95,96,97}

7. Acupuncture is widely available from qualified practitioners nationally.

In 2013 more than 28,000 licensed acupuncturists were estimated to be practicing in the U.S., with many more in training.⁹⁸ A 2015 study found the number of trained practitioners to be approximately 34,400. The number of licensed acupuncturists was noted to have increased by 23.3% and 52.1% compared to the years 2009 (n=27,965) and 2004 (n=22,671) respectively.⁹⁹ Currently the Council of Colleges of Acupuncture and Oriental Medicine (CCAOM) has 57 schools in its membership,¹⁰⁰ with approximately 10 schools offering doctoral degrees. The National Certification Council for Acupuncture and Oriental Medicine (NCCAOM) has certified more than 18,000 practitioners for minimal competency.¹⁰¹ The American Academy of Medical Acupuncture

also represents more than 1,300 medical doctors trained to offer acupuncture services, and has approved nine programs for medical doctor certification in acupuncture.¹⁰² One certification program alone has trained more than 6000 physicians in medical acupuncture,¹⁰³ so a conservative estimate of the total number of physicians trained would be approximately 10,000, though the number in active practice with acupuncture is unknown. Most states allow physicians to practice acupuncture, with some specifying additional training.¹⁰⁴ Increased coverage and demand for acupuncture will lead to a greater supply of providers as well. As noted above, NADA providers are estimated at 25,000 individuals, with more than 1,000 programs in the U.S. and Canada.

Appendix 1. Effectiveness of Acupuncture.

Author, Year	Topic/Intervention	Participants/Population	Primary Outcomes	Key Findings	Study Quality
Vickers et al, 2012	Acupuncture versus sham acupuncture and no acupuncture in back, neck, shoulder pain, chronic headache, osteoarthritis	Systematic review of 31 randomized controlled trials (17,922 subjects) and meta-analysis of individual patient data from 29 of these 31 RCTs in back, neck, shoulder pain; chronic headache, osteoarthritis	A variety of pain severity and disability scores such as VAS, WOMAC, Roland Morris Disability Questionnaire	Acupuncture was superior to sham acupuncture and no acupuncture for each pain condition	High quality evidence
Weidenhammer et al, 2007	Acupuncture for headache, low back pain, osteoarthritis	Open pragmatic trial of 454,920 subjects with headache, low back pain, osteoarthritis	Treating physician rating of “marked, moderate, minimal or poor improvement (which included no improvement and worse)”	Physician ratings: 22% marked, 54% moderate, 16% minimal and 4% poor improvement	Low quality evidence - Open pragmatic trial with no blinding and no external assessors
Corbett et al, 2013	Comparison of 22 physical therapies for knee osteoarthritis pain	Review of 152 trials and network meta-analysis of 12 randomized controlled trials with low risk of bias comparing 22 physical therapies in knee osteoarthritis pain	Knee pain	Acupuncture was equal to balneotherapy and superior to sham acupuncture, muscle-strengthening exercise, tai chi, weight loss, standard care and aerobic exercise (in ranked order)	110 of 152 studies analysed were of poor quality. Network meta-analysis included 12 RCTs with low risk of bias

Ji et al, 2015	Acupuncture versus standard pharmaceutical care in sciatica	Systematic review and meta-analysis of 12 randomized controlled trials in sciatica	Effectiveness, pain intensity, pain threshold	Acupuncture was superior to standard pharmaceutical care in effectiveness, reducing pain intensity and pain threshold	Low to moderate quality evidence
Lewis et al, 2015	Comparison of 21 different interventions for sciatica	Systematic review and network meta-analyses of 122 studies including 90 randomized or quasi-randomized controlled trials comparing 21 different interventions for sciatica	Global effect, pain intensity	In global effect and reduction in pain intensity, acupuncture was second only to biological agents (cytokine modulating drugs), and superior to all other interventions tested including non-opioid and opioid medications	9% of studies had a strong overall quality rating; 7% of studies had a strong overall external validity rating; 21% of studies used both adequate randomization and adequate or partially adequate allocation concealment
Gadau et al, 2014	Acupuncture and/or moxibustion versus sham acupuncture, another form of acupuncture, or conventional treatment in lateral elbow pain	Systematic review of 19 randomized controlled trials	Pain, grip strength	Acupuncture is more effective than sham acupuncture (moderate quality studies) Acupuncture or moxibustion is more effective than conventional treatment (low quality studies)	Low to moderate quality evidence

Cho et al, 2015	Real versus sham acupuncture in acute post-operative pain after back surgery	Systematic review and meta-analysis of 5 trials	24-hour post-operative pain intensity on VAS; 24-hour opiate demands	Real acupuncture was superior to sham in reducing pain intensity but not opiate demand at 24-hours	3 of 5 trials were high quality
Levett et al, 2014	Acupuncture, standard care, sham acupuncture, acupressure and mixed controls in various combinations in labor pain	A critical narrative review of 4 systematic reviews in labor pain	Pain intensity, analgesic use, length of labor	Acupuncture reduces pain intensity, analgesic use and length of labor	Conflicting evidence
Clark et al, 2012	Acupuncture versus various comparators including standard care, sham acupuncture and other forms of acupuncture in plantar heel pain	Systematic review of 5 randomized controlled trials and 3 non-randomized comparative trials	Various pain and disability scales (morning pain, walking pain, tenderness)	Acupuncture for plantar heel pain is supported by evidence which is equivalent to evidence supporting standard care (stretching, splints, dexamethasone)	Evidence at level I and II supporting the effectiveness of acupuncture for heel pain, leading to a recommendation at Grade B
Deare et al, 2014	Manual and electro-acupuncture compared with sham acupuncture, standard therapy and no treatment in fibromyalgia	Cochrane systematic review of 9 randomized controlled trials in fibromyalgia	Pain, stiffness, sleep, fatigue and global wellbeing	Acupuncture improves pain and stiffness compared to standard therapy and no treatment, but not compared to sham acupuncture	Low to moderate quality evidence

Smith et al, 2011	Acupuncture or acupressure versus placebo control, usual care or pharmacological treatment in primary dysmenorrhea	Cochrane systematic review of 10 randomized controlled trials (944 subjects) on acupuncture (6) or acupressure (4) for primary dysmenorrhea	Pain relief, analgesic use, quality of life, improvement in menstrual symptoms, absenteeism	Acupuncture was superior to placebo and Chinese herbs in pain relief, and superior to medication and Chinese herbs in reducing menstrual symptoms. Acupressure was superior to placebo in pain relief and reducing menstrual symptoms	Low risk of bias in 50% of included RCTs
Abaraogu et al, 2015	Acupuncture or acupressure versus placebo control, wait list or pharmacological treatment in primary dysmenorrhea	Systematic review of 8 randomized controlled trials (>3,000 subjects) and meta-analysis of 4 RCTs	Pain intensity (VAS, McGill), quality of life, blood nitric oxide	Acupuncture and acupressure reduced pain, while acupuncture also improved quality of life	Moderate quality evidence
Chen et al, 2013	Acupuncture or acupressure at acupoint SP 6 versus minimal stimulation at SP 6 or stimulation of another point in primary dysmenorrhea	Meta-analysis of acupuncture (3) and acupressure (4) randomized controlled trials in primary dysmenorrhea	Pain intensity (VAS)	Acupuncture is effective and acupressure may be effective at SP 6 for pain relief	Acupuncture trials had low to moderate risk of bias Acupressure trials had high risk of bias

Cho et al, 2010	Acupuncture versus sham acupuncture, pharmacological treatment or Chinese herbs in primary dysmenorrhea	Systematic review of 27 randomized controlled trials in primary dysmenorrhea	Pain intensity (VAS, Menstrual Pain Reduction Score, other pain scores)	Acupuncture was superior to pharmacological treatment or Chinese herbs in pain relief	Only 5 out of 27 trials had low risk of bias
Chung et al, 2012	Acupoint stimulation versus non-acupoint stimulation or medication in primary dysmenorrhea	Systematic review of 30 randomized controlled trials (>3,000 subjects) and meta-analysis of 25 RCTs	Pain intensity, plasma PGF(2)/PGE(2) ratio	Acupoint stimulation was superior in short-term pain relief to stimulation on non-acupoints. Non-invasive stimulation of acupoints was more effective than invasive stimulation	Some trials were of low quality
Xu et al, 2014	Various forms of acupoint stimulation (including acupuncture, moxibustion and other methods) versus a variety of controls in primary dysmenorrhea	Meta-analysis of 20 randomized controlled trials (2,134 subjects) of acupoint stimulation for primary dysmenorrhea	Pain relief	Acupoint stimulation was more effective than controls for pain relief	Low to moderate quality evidence

References

- ¹ McDonald J, Janz S. The Acupuncture Evidence Project: A Comprehensive Literature Review. Australian Acupuncture & Chinese Medicine Association Limited, Dec 19, 2016
- ² Vickers AJ, Cronin AM, Maschino AC, Lewith G, MacPherson H, Foster NE, et al. Acupuncture for Chronic Pain: Individual Patient Data Meta-Analysis. *Archives of Internal Medicine*. 2012;172(19):1444-53.
- ³ Weidenhammer W, Streng A, Linde K, Hoppe A, Melchart D. Acupuncture for Chronic Pain within the Research Program of 10 German Health Insurance Funds--Basic Results from an Observational Study. *Complementary Therapies in Medicine*. 2007;15(4):238-46.
- ⁴ Corbett MS, Rice SJ, Madurasinghe V, Slack R, Fayter DA, Harden M, et al. Acupuncture and other Physical Treatments for the Relief of Pain Due to Osteoarthritis of the Knee: Network Meta-analysis. *Osteoarthritis and Cartilage / OARS, Osteoarthritis Research Society*. 2013;21(9):1290-8.
- ⁵ Qaseem A, Wilt TJ, McLean RM, Forcica MA; Clinical Guidelines Committee of the American College of Physicians. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline from the American College of Physicians. *Ann Intern Med*. 2017 Apr 4;166(7):514-530. doi: 10.7326/M16-2367. Epub 2017 Feb 14.
- ⁶ Ji M, Wang X, Chen M, Shen Y, Zhang X, Yang J. The Efficacy of Acupuncture for the Treatment of Sciatica: A Systematic Review and Meta-Analysis. *Evidence-based Complementary and Alternative Medicine: eCAM*. 2015;2015:192808.
- ⁷ Lewis RA, Williams NH, Sutton AJ, Burton K, Din NU, Matar HE, et al. Comparative Clinical Effectiveness of Management Strategies for Sciatica: Systematic Review and Network Meta-analyses. *The Spine Journal: Official Journal of the North American Spine Society*. 2015;15(6):1461-77.
- ⁸ Gadau M, Yeung WF, Liu H, Zaslowski C, Tan YS, Wang FC, et al. Acupuncture and Moxibustion for Lateral Elbow Pain: A Systematic Review of Randomized Controlled Trials. *BMC Complementary and Alternative Medicine*. 2014;14:136.
- ⁹ Clark RJ, Tighe M. The Effectiveness of Acupuncture for Plantar Heel Pain: A Systematic Review. *Acupuncture in Medicine: Journal of the British Medical Acupuncture Society*. 2012;30(4):298-306.
- ¹⁰ An LX, Chen X, Ren XJ, Wu HF. Electro-Acupuncture Decreases Postoperative Pain and Improves Recovery in Patients Undergoing Supratentorial Craniotomy. *The American Journal of Chinese Medicine*. 2014;42(5):1099-109.
- ¹¹ Chen CC, Yang CC, Hu CC, Shih HN, Chang YH, Hsieh PH. Acupuncture for Pain Relief after Total Knee Arthroplasty: A Randomized Controlled Trial. *Regional Anesthesia and Pain Medicine*. 2015;40(1):31-6.
- ¹² Cho HK, Park IJ, Jeong YM, Lee YJ, Hwang SH. Can Perioperative Acupuncture Reduce the Pain and Vomiting Experienced after Tonsillectomy? A Meta-Analysis. *The Laryngoscope*. 2015.
- ¹³ Cho YH, Kim CK, Heo KH, Lee MS, Ha IH, Son DW, et al. Acupuncture for Acute Postoperative Pain after Back Surgery: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Pain Practice: The Official Journal of World Institute of Pain*. 2015;15(3):279-91.
- ¹⁴ Crespin DJ, Griffin KH, Johnson JR, Miller C, Finch MD, Rivard RL, et al. Acupuncture Provides Short-Term Pain Relief for Patients in a Total Joint Replacement Program. *Pain Medicine (Malden, Mass)*. 2015;16(6):1195-203.
- ¹⁵ Gilbey P, Bretler S, Avraham Y, Sharabi-Nov A, Ibrgimov S, Luder A. Acupuncture for Post Tonsillectomy Pain in Children: A Randomized, Controlled Study. *Paediatric Anaesthesia*. 2015;25(6):603-9.
- ¹⁶ Liu XL, Tan JY, Molassiotis A, Suen LK, Shi Y. Acupuncture-Point Stimulation for Postoperative Pain Control: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Evidence-based complementary and alternative medicine: eCAM*. 2015;2015:657809.
- ¹⁷ Lu Z, Dong H, Wang Q, Xiong L. Perioperative Acupuncture Modulation: More than Anaesthesia. *British Journal of Anaesthesia*. 2015;115(2):183-93.
- ¹⁸ Tsao GJ, Messner AH, Seybold J, Sayyid ZN, Cheng AG, Golianu B. Intraoperative Acupuncture for Post Tonsillectomy Pain: A Randomized, Double-Blind, Placebo-Controlled Trial. *The Laryngoscope*. 2015;125(8):1972-8.
- ¹⁹ Ibid, Cho HK, Park IJ, et al.
- ²⁰ Ibid, Cho HK, Kim CK, et al.
- ²¹ Golianu B, Krane E, Seybold J, Almgren C, Anand KJS. Non-Pharmacological Techniques for Pain Management in Neonates. *Seminars in Perinatology*. 2007;31(5):318-22.
- ²² Fry LM, Neary SM, Sharrock J, Rychel JK. Acupuncture for Analgesia in Veterinary Medicine. *Topics in Companion Animal Medicine*. 2014;29(2):35-42.

- ²³ Levy B, Paulozzi L, Mack KA, Jones CM. Trends in Opioid Analgesic-Prescribing Rates by Specialty, U.S., 2007-2012. *American Journal of Preventive Medicine*. 2015; 49(3), 409-413. doi:10.1016/j.amepre.2015.02.020
- ²⁴ Thiels CA, Anderson SS, Ubl DS, et al. Wide Variation and Overprescription of Opioids after Elective Surgery. *Annals of Surgery*. 2017: doi:10.1097/sla.0000000000002365
- ²⁵ Wunsch H, Wijeyesundera DN, Passarella MA, et al. Opioids Prescribed after Low-Risk Surgical Procedures in the United States, 2004-2012. *JAMA*. 2016;315(15), 1654-1657. doi:10.1001/jama.2016.0130
- ²⁶ Hill MV, McMahon ML, Stucke RS, Barth RJ. Wide Variation and Excessive Dosage of Opioid Prescriptions for Common General Surgical Procedures. *Annals of Surgery*. 2017: 265(4), 709-714. doi:10.1097/sla.0000000000001993
- ²⁷ Brummett CM, Waljee JF, Goesling J, Moser S, Lin P, et al. New Persistent Opioid Use after Minor and Major Surgical Procedures in US Adults. *JAMA Surg*, 2017;152(6), e170504. doi:10.1001/jamasurg.2017.0504
- ²⁸ Pang J, Tringale KR, Tapia VJ, Moss WJ, et al (2017). Chronic Opioid Use Following Surgery for Oral Cavity Cancer. *JAMA Otolaryngol Head Neck Surgery*. 2017: doi:10.1001/jamaoto.2017.0582
- ²⁹ Meissner W, Dohrn B, Reinhart K (2003). Enteral Naloxone Reduces Gastric Tube Reflux and Frequency of Pneumonia in Critical Care Patients During Opioid Analgesia. *Critical Care Medicine*. 2003;31(3), 776-780. doi:10.1097/01.ccm.0000053652.80849.9f
- ³⁰ Dublin S, Walker RL, Jackson ML, et al. Use of Opioids or Benzodiazepines and Risk of Pneumonia In Older Adults: A Population-Based Case-Control Study. *Journal of the American Geriatrics Society*. 2011;59(10), 1899-1907. doi:10.1111/j.1532-5415.2011.03586.x
- ³¹ Burry LD, Williamson DR, Mehta S, Perreault MM, et al. Delirium and Exposure to Psychoactive Medications in Critically Ill Adults: A Multi-centre Observational Study. *Journal of Critical Care*. 2017: 42, 268-274. doi:10.1016/j.jcrc.2017.08.003
- ³² Tedesco D, Gori D, Desai KR, Asch S, Carroll IR, Curtin C, McDonald KM, Fantini MP, Hernandez-Boussard T. Drug-Free Interventions to Reduce Pain or Opioid Consumption After Total Knee Arthroplasty: A Systematic Review and Meta-analysis. *JAMA Surg*. 2017 Aug 16:e172872. doi: 10.1001/jamasurg.2017.2872. [Epub ahead of print]
- ³³ Asmussen S, Przkora R, Maybauer DM, Fraser JF, Sanfilippo F, Jennings K, Maybauer MO. Meta-Analysis of Electroacupuncture in Cardiac Anesthesia and Intensive Care. *Journal of Intensive Care Medicine*. 2017 885066617708558. doi:10.1177/0885066617708558
- ³⁴ Huang S, Peng, W, Tian X, et al. Effects of Transcutaneous Electrical Acupoint Stimulation at Different Frequencies on Perioperative Anesthetic Dosage, Recovery, Complications, and Prognosis in Video-Assisted Thoracic Surgical Lobectomy: A Randomized, Double-Blinded, Placebo-Controlled Trial. *Journal of Anesthesia*. 2017;31(1), 58-65. doi:10.1007/s00540-015-2057-1
- ³⁵ An LX, Chen X, Ren XJ, & Wu HF. Electro-Acupuncture Decreases Postoperative Pain and Improves Recovery in Patients Undergoing a Supratentorial Craniotomy. *American Journal of Chinese Medicine*. 2014;42(5), 1099-1109. doi:10.1142/s0192415x14500682
- ³⁶ Asmussen S, Maybauer, DM, Chen JD, Fraser JF, Toon MH, Przkora R, Maybauer MO. Effects of Acupuncture in Anesthesia for Craniotomy: A Meta-Analysis. *Journal of Neurosurgical Anesthesiology*. 2017;29(3), 219-227. doi:10.1097/ana.0000000000000290
- ³⁷ Yang Y, Zuo HQ, Li Z, et al. Comparison of Efficacy of Simo Decoction and Acupuncture or Chewing Gum Alone on Postoperative Ileus in Colorectal Cancer Resection: A Randomized Trial. *Scientific Reports*. 2017: 7, 37826. doi:10.1038/srep37826
- ³⁸ Smith CA, Zhu X, He L, Song J. Acupuncture for Primary Dysmenorrhoea. The Cochrane Database of Systematic Reviews. 2011(1): Cd007854.
- ³⁹ Abaraogu UO, Tabansi-Ochuogu CS. As Acupressure Decreases Pain, Acupuncture may Improve Some Aspects of Quality of Life for Women with Primary Dysmenorrhea: A Systematic Review with Meta-Analysis. *Journal of Acupuncture and Meridian Studies*. 2015;8(5):220-8.
- ⁴⁰ Cho SH, Hwang EW. Acupuncture for Primary Dysmenorrhoea: A Systematic Review. *BJOG : An International Journal of Obstetrics and Gynaecology*. 2010;117(5):509-21.
- ⁴¹ Xu T, Hui L, Juan YL, Min SG, Hua WT. Effects of Moxibustion or Acupoint Therapy for the Treatment of Primary Dysmenorrhea: A Meta-Analysis. *Alternative Therapies in Health and Medicine*. 2014;20(4):33-42.
- ⁴² Chung YC, Chen HH, Yeh ML. Acupoint Stimulation Intervention for People with Primary Dysmenorrhea: Systematic Review and Meta-Analysis of Randomized Trials. *Complementary Therapies in Medicine*. 2012;20(5):353-63.
- ⁴³ Chen MN, Chien LW, Liu CF. Acupuncture or Acupressure at the Sanyinjiao (SP6) Acupoint for the Treatment of Primary Dysmenorrhea: A Meta-Analysis. *Evidence-based Complementary and Alternative Medicine : eCAM*. 2013;2013:493038.

-
- ⁴⁴ Levett KM, Smith CA, Dahlen HG, Bensoussan A. Acupuncture and Acupressure for Pain Management in Labour and Birth: A Critical Narrative Review of Current Systematic Review Evidence. *Complementary Therapies in Medicine*. 2014;22(3):523-40.
- ⁴⁵ Vixner L, Schytt E, Stener-Victorin E, Waldenstrom U, Pettersson H, Martensson LB. Acupuncture with Manual and Electrical Stimulation for Labour Pain: A Longitudinal Randomised Controlled Trial. *BMC Complementary and Alternative Medicine*. 2014;14:187.
- ⁴⁶ Dong C, Hu L, Liang F, Zhang S. Effects of Electro-Acupuncture on Labor Pain Management. *Archives of Gynecology and Obstetrics*. 2015;291(3):531-6.
- ⁴⁷ Liu H, Li H, Xu M, Chung KF, Zhang SP. A Systematic Review on Acupuncture for Trigeminal Neuralgia. *Alternative Therapies in Health and Medicine*. 2010;16(6):30-5.
- ⁴⁸ Deare JC, Zheng Z, Xue CC, Liu JP, Shang J, Scott SW, et al. Acupuncture for Treating Fibromyalgia. *The Cochrane Database of Systematic Reviews*. 2013;5:CD007070.
- ⁴⁹ Grissa MH, Baccouche H, Boubaker H, Beltaief K, Bzeouich N, Fredj N, et al. Acupuncture vs Intravenous Morphine in the Management of Acute Pain in the ED. *American Journal of Emergency Med*. 2016;34(11):2112-2116.
- ⁵⁰ Ibid, Vickers AJ, Cronin AM, et al.
- ⁵¹ Ibid, Weidenhammer W, Streng A, et al.
- ⁵² Feeney C, Bruns E, LeCompte G, Forati A, Chen T, Matecki A. Acupuncture for Pain and Nausea in the Intensive Care Unit: A Feasibility Study in a Public Safety Net Hospital, *Journal of Alternative and Complementary Medicine*. 2017 Apr 25. doi: 10.1089/acm.2016.0323. [Epub ahead of print]
- ⁵³ Lao L. Acupuncture Practice, Past and Present: Is it Safe and Effective? *Journal of the Society of Integrative Oncology*. 2006;4(1):13-5.
- ⁵⁴ Lu W, Dean-Clower E, et al. The Value of Acupuncture in Cancer Care. *Hematol Oncol Clin North Am*. 2008: Aug; 22(4): 631–viii. doi: 10.1016/j.hoc.2008.04.005
- ⁵⁵ Ambrosio EM, Bloor K, MacPherson H. Costs and Consequences of Acupuncture as a Treatment for Chronic Pain: A Systematic Review of Economic Evaluations Conducted Alongside Randomised Controlled Trials. *Complementary Therapies in Medicine*. 2012;20(5):364-74.
- ⁵⁶ MacPherson H, Vickers A, Bland JM, Torgerson DJ, Corbett MS, Spackman E, Saramago Goncalves PR, Woods BS, Weatherly HL, Sculpher MJ, Manca A. Acupuncture for Chronic Pain and Depression in Primary Care: A Programme of Research. *Programme Grants for Applied Research*. 2017 Jan 1:1-342.
- ⁵⁷ CHIA center for health information and analysis. Mandated Benefit Review Of H.B. 3972: An Act Relative to the Practice of Acupuncture. <http://www.aomsm.org/Resources/Documents/Research/BenefitReview-H3972-Acupuncture.pdf> Accessed August 19, 2017.
- ⁵⁸ http://www.acupuncture.org.au/Portals/0/Evidence%20study/Acupuncture%20Evidence_plain%20English%20Web%20version_17_Feb.pdf?ver=2017-02-22-171448-550. Accessed 8/26/17.
- ⁵⁹ Lin JG, Lo MW, Wen YR, Hsieh CL, Tsai SK, Sun WZ. The Effect of High and Low Frequency Electroacupuncture in Pain after Lower Abdominal Surgery. *Pain*. 2002;99(3):509-14.
- ⁶⁰ Wang B, Tang J, White PF, Naruse R, Sloninsky A, Kariger R, et al. Effect of the Intensity of Transcutaneous Acupoint Electrical Stimulation on the Postoperative Analgesic Requirement. *Anesthesia and Analgesia*. 1997;85(2):406-13.
- ⁶¹ Zheng Z, Guo RJ, Helme RD, Muir A, Da Costa C, Xue CC. The Effect of Electroacupuncture on Opioid-Like Medication Consumption by Chronic Pain Patients: A Pilot Randomized Controlled Clinical Trial. *European Journal of Pain (London, England)*. 2008;12(5):671-6.
- ⁶² Ibid, Tedesco D, Gori D, et al.
- ⁶³ Crawford Paul, Penzien Donald B., and Coeytaux Remy. *Medical Acupuncture*. August 2017, 29(4): 229-231. <https://doi.org/10.1089/acu.2017.1234>
- ⁶⁴ Kligler, B. Integrative Health in the Veterans Health Administration. *Medical Acupuncture*. 2017;29(4):187-188. doi: 10.1089/acu.2017.29055.bkl
- ⁶⁵ Helms, J. Medical Acupuncture Meets the Military. *Medical Acupuncture*. 2017;29(4):189-190. doi: 10.1089/acu.2017.29055.bkl
- ⁶⁶ Hanlon JT, Zhao X, et al. Central Nervous System Medication Burden and Serious Falls in Older Nursing Home Residents. *J Am Geriatr Soc*. 2017 Jun;65(6):1183-1189. doi: 10.1111/jgs.14759. Epub 2017 Feb 2.
- ⁶⁷ Robb G, Loe E, et al. Medication-Related Patient Harm in New Zealand Hospitals. *N Z Med J*. 2017 Aug 11;130(1460):21-32.

-
- ⁶⁸ Goldman N, Chen M, Fujita T, Xu Q, Peng W, Liu W, Jensen TK, Pei Y, Wang F, Han X, Chen JF. Adenosine A1 Receptors Mediate Local Anti-Nociceptive Effects of Acupuncture. *Nature Neuroscience*. 2010 Jul 1;13(7):883-8.
- ⁶⁹ Takano T, Chen X, Luo F, Fujita T, Ren Z, Goldman N, Zhao Y, Markman JD, Nedergaard M. Traditional Acupuncture Triggers a Local Increase in Adenosine in Human Subjects. *The Journal of Pain*. 2012 Dec 31;13(12):1215-23.
- ⁷⁰ Zhao ZQ. Neural Mechanism Underlying Acupuncture Analgesia. *Progress in Neurobiology*. 2008;85(4):355-75.
- ⁷¹ Han JS. Acupuncture Analgesia: Areas of Consensus and Controversy. *Pain*. 2011;152(3 Suppl):S41-8.
- ⁷² Han JS. Acupuncture and Endorphins. *Neuroscience Letters*. 2004;361(1-3):258-61.
- ⁷³ McDonald JL, Cripps AW, Smith PK. Mediators, Receptors, and Signalling Pathways in the Anti-Inflammatory and Antihyperalgesic Effects of Acupuncture. *Evidence-based Complementary and Alternative Medicine : eCAM*. 2015;2015:975632.
- ⁷⁴ Zhang R, Lao L, Ren K, Berman BM. Mechanisms of Acupuncture-Electroacupuncture on Persistent Pain. *Anesthesiology*. 2014;120(2):482-503.
- ⁷⁵ Harris RE, Zubieta JK, Scott DJ, Napadow V, Gracely RH, Clauw DJ. Traditional Chinese Acupuncture and Placebo (sham) Acupuncture are Differentiated by Their Effects on μ -opioid Receptors (MORs). *Neuroimage*. 2009 Sep 30;47(3):1077-85.
- ⁷⁶ Ibid, Qaseem A, et al.
- ⁷⁷ Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain — United States, 2016. *MMWR Recomm Rep* 2016;65(No. RR-1):1–49. doi: <http://dx.doi.org/10.15585/mmwr.rr6501e1>.
- ⁷⁸ Shaheed CA, Maher CG, Williams KA, Day R, McLachlan AJ. Efficacy, Tolerability, and Dose-dependent Effects of Opioid Analgesics for Low Back Pain: A Systematic Review and Meta-analysis. *JAMA Internal Medicine*. 2016 Jul 1;176(7):958-68.
- ⁷⁹ Krebs EE. Effectiveness of Opioid Therapy Vs. Non-Opioid Medication Therapy for Chronic Back and Osteoarthritis Pain Over 12 Months. In Annual Meeting, Society for General Internal Medicine, Washington DC 2017.
- ⁸⁰ Xing GG, Liu FY, Qu XX, Han JS, Wan Y. Long-Term Synaptic Plasticity in the Spinal Dorsal Horn and its Modulation by Electroacupuncture in Rats with Neuropathic Pain. *Experimental Neurology*. 2007;208(2):323-32.
- ⁸¹ Napadow V, Kettner N, Ryan A, Kwong KK, Audette J, Hui KK. Somatosensory Cortical Plasticity in Carpal Tunnel Syndrome--A Cross-Sectional Fmri Evaluation. *NeuroImage*. 2006;31(2):520-30.
- ⁸² Napadow V, Liu J, Li M, Kettner N, Ryan A, Kwong KK, et al. Somatosensory Cortical Plasticity in Carpal Tunnel Syndrome Treated by Acupuncture. *Human Brain Mapping*. 2007;28(3):159-71.
- ⁸³ Liu CZ, Kong J, Wang KL. Acupuncture Therapies and Neuroplasticity. *Neural Plast*. 2017; 2017: 6178505. Published online 2017 Apr 27. doi: 10.1155/2017/6178505
- ⁸⁴ US National Library of Medicine National Institutes of Health, Search Database. <https://www.ncbi.nlm.nih.gov/pubmed/?term=acupuncture+opioids+addition> Accessed August 18, 2017.
- ⁸⁵ Wen H, Cheung SYC. Treatment of Drug Addiction by Acupuncture and Electrical Stimulation. *Asian J Med*. 1973;9:138-41.
- ⁸⁶ National Acupuncture Detoxification Association. About NADA. <http://www.acudetox.com/about-nada/12-faqs2013>.
- ⁸⁷ Lin JG, Chan YY, Chen YH. Acupuncture for the Treatment of Opiate Addiction. *Evidence Based Complement Alternative Med*. 2012;2012:739045.
- ⁸⁸ Raith W, Schmolzer GM, Resch B, Reiterer F, Avian A, Koestenberger M, et al. Laser Acupuncture for Neonatal Abstinence Syndrome: A Randomized Controlled Trial. *Pediatrics*. 2015;136(5):876-84.
- ⁸⁹ Wu LZ, Cui CL, Tian JB, Ji D, Han JS. Suppression of Morphine Withdrawal by Electroacupuncture in Rats: Dynorphin and Kappa-Opioid Receptor Implicated. *Brain Research*. 1999;851(1-2):290-6.
- ⁹⁰ The Food and Drug Administration. Introduction for the FDA Blueprint for Prescriber Education for Extended-Release and Long-Acting Opioid Analgesics. <https://www.fda.gov/downloads/Drugs/DrugSafety/InformationbyDrugClass/UCM515636.pdf> Accessed August 19, 2017.
- ⁹¹ The National Academies of Science, Engineering and Medicine. National Strategy to Reduce Opioid Epidemic, an Urgent Public Health Priority, Presented in New Report. <http://www8.nationalacademies.org/onpinews/newsitem.aspx?RecordID=24781> Accessed August 19, 2017; <https://www.nap.edu/read/24781/chapter/4>, pp. 69-70.

-
- ⁹² Official Publication of Joint Commission Requirements New and Revised Standards Related to Pain Assessment and Management, Volume 37, Number 7, July 2017.
https://www.jointcommission.org/assets/1/18/Joint_Commission_Enhances_Pain_Assessment_and_Management_Requirements_for_Accredited_Hospitals1.PDF.
- ⁹³ Birch S, Alraek T, Lee MS. Challenges for Clinical Practice Guidelines in Traditional Medicines: The Example of Acupuncture. *European Journal of Integrative Medicine*. 2016 Aug 31;8(4):332-6.
- ⁹⁴ Ibid, Qaseem A, et al.
- ⁹⁵ Ibid, Todesco D, et al.
- ⁹⁶ Ibid, Grissa MH, et al.
- ⁹⁷ Ibid, CHIA center Mandated Benefit Review.
- ⁹⁸ Academic Collaborative for Integrative Health (ACIH), Clinicians' & Educators' Desk Reference on the Integrative Health & Medicine Professions Third Edition, Mercer Island, WA, 2017, Pp. 17-42.
https://static1.squarespace.com/static/55861f1ae4b01ea9a58583a7/t/597bb31e914e6b7dd5beb083/1501279018792/2017+CEDR_final_071817.pdf. Accessed 8/26/17.
- ⁹⁹ Fan AY, Faggert S. Number of Licensed Acupuncturists and Educational Institutions in the United States in Early of 2015. *J Integrat Med*. 2017 September; Epub ahead of print. doi:10.1016/S2095-4964(17)60371-6
- ¹⁰⁰ www.ccaom.org
- ¹⁰¹ www.nccaom.org
- ¹⁰² <http://www.medicalacupuncture.org/>
- ¹⁰³ <https://hmieducation.com/about-hmi>
- ¹⁰⁴ Lin K, Tung, C. The Regulation of the Practice of Acupuncture by Physicians in the United States. *Medical Acupuncture*. 2017; 29(3): 121-125.